arXiv:cond-mat/0401597v3 [cond-mat.mes-hall] 3 Mar 2004

Fractional flux periodicity of a twisted planar square lattice

K. Sasaki* and Y. Kawazoe
Institute for Materials Research, Tohoku University, Sendai 980-8577, Japan

R. Saito
Department of Physics, Tohoku University and CREST, JST, Sendai 980-8578, Japan

(Dated: February 2, 2008)

We present fractional flux periodicity in the ground state of planar systems made of a square lattice whose boundary is compacted into a torus. The ground-state energy shows a fractional period of the fundamental unit of magnetic flux depending on the twist around the torus axis.

The Aharonov-Bohm effect [1] shows that a single electron wave function has a fundamental unit of magnetic flux $\Phi_0 = 2\pi/e$, where $-e$ is the electron charge. We will use the units of $\hbar = c = 1$. The electric and magnetic properties of materials are governed by many electrons, and each constituent has the above-mentioned periodicity. However, the fundamental flux period of a material need not be Φ_0. For example, superconducting materials exhibit a period one half the single-electron flux quantum, that is, $\Phi_0/2$, which can be understood by charge doubling due to the Cooper pair formation. This is clear if one imagine that a fundamental particle (or quasiparticle) has a $-2e$ charge due to the attractive interaction and that the fundamental flux period of a material is equal to that of the quasiparticle ($2\pi/2e$) in the system. So, one may ask: Is it possible that fractional flux periodicity ($\Phi_0/2, \Phi_0/3, \cdots$) realizes in the absence of the interaction? We answer this question for torus geometries whose surface is a planar system made of a square lattice. To analyze the flux periodicity of the systems, we will consider the persistent currents [2, 3], which are suitable to examine the flux periodicity of the ground state since they are regarded as the Aharonov-Bohm effect in solid state systems [4].

The quantum mechanical behavior of conducting electrons on a square lattice is modeled by the nearest-neighbor tight-binding Hamiltonian:

$$\mathcal{H} = -t \sum_i \sum_{a=x,y} a^\dagger_{i+a} e^{-ieA^{\rm ex}\cdot T_a} a_i + h.c., \tag{1}$$

where t is the hopping integral, $A^{\rm ex}$ is a constant external gauge field (vector potential), and T_a ($a = x, y$) is the vector connecting each site in the direction of x and y. a_i and $a^\dagger_i$ are canonical annihilation-creation operators of the electrons at site i that satisfy the anti-commutation relation $\{a_i, a^\dagger_j\} = \delta_{ij}$. The energy eigenvalue of the Hamiltonian is parametrized by the Bloch wave vector k as

$$E(k - eA^{\rm ex}) = -2t\Re\left[\sum_{a=x,y} e^{i(k-eA^{\rm ex})\cdot T_a}\right]. \tag{2}$$

The geometry of a torus can be specified by two vectors: chiral $C_h = NT_x + MT_y$ and translational $T_w = PT_x + QT_y$ vectors (we borrow this terminology from the carbon nanotube context [5]), where N,M,P, and Q are integers. It is useful to rewrite T_a in terms of the chiral and translational vectors as

$$\begin{pmatrix} T_x \\ T_y \end{pmatrix} = \frac{1}{N_s}\begin{pmatrix} Q & -M \\ -P & N \end{pmatrix}\begin{pmatrix} C_h \\ T_w \end{pmatrix}, \tag{3}$$

where we define $N_s = NQ - MP$. In this study we will take x axis in the direction of C_h ($M = 0$), and denote $P = \delta N$ to describe the *twist* around the tube axis. Then, $C_h \cdot T_w = \delta N N a^2$ where $a(= |T_x| = |T_y|)$ is the lattice spacing. Figure 1(a) illustrates a twisted torus. It should be noted that a torus can be unrolled to a parallelogram sheet as depicted in Fig. 1(b). In the figure, the two lines extending upward from 'u' and downward from 'd' and having the same 'x' at the junction are not joined for a twisted torus, shown in the lower inset of Fig. 1(a). There are δN square lattices between the two lines.


(a)
d
u
d
δN
u
(c)
(b)
u
$A^{\rm ex}$
T_w
$\frac{A^{\rm ex}\cdot T_w}{|T_w|}$
$\frac{A^{\rm ex}\cdot C_h}{|C_h|}$
d
y
x
C_h

FIG. 1: Twisted torus (a), its net diagram (b) with an external gauge field, and an example of a *twisted* torus (c). It is convenient to consider a parallelogram as the net of a twisted torus. We draw several square lattices in (b).

The Bloch wave vector satisfies the periodic boundary condition through which the geometrical (or topological) information, such as the twist, is put into the energy eigenvalue. We decompose wave vector k as $\mu_1 k_1 + \mu_2 k_2$,

---

*Electronic address: sasaken@imr.edu

where μ_1 and μ_2 are integers. Here, k_1 and k_2 are defined by

$$C_h \cdot k_1 = 2\pi, \; C_h \cdot k_2 = 0, \; T_w \cdot k_1 = 0, \; T_w \cdot k_2 = 2\pi. \tag{4}$$

By means of Eqs.(3) (with $M = 0$ and $P = \delta N$) and (4), we rewrite Eq.(2) as

$$E(k - eA^{\rm ex}) = -2t \, \Re \left[e^{i \frac{2\pi\mu_1}{N}} + e^{i \frac{2\pi(\mu_2 - \frac{\delta N}{N}\mu_1) - eA^{\rm ex}\cdot T_w}{Q}} \right], \tag{5}$$

where we assume $A^{\rm ex} \cdot C_h = 0$, and $A^{\rm ex} \cdot T_w$ corresponds to the Aharonov-Bohm flux penetrating the center of the torus. It is important to note that the twist works as an extra gauge field, $A^{\rm twist}$, along the axis [6], because the second term of the right hand side of Eq.(5) can be rewritten as $\exp\left(i \frac{2\pi\mu_2 - (eA^{\rm ex} + \mu_1 A^{\rm twist})\cdot T_w}{Q}\right)$, where we define $A^{\rm twist} \cdot C_h \equiv 0$ and $A^{\rm twist} \cdot T_w \equiv 2\pi\delta N/N$. Like $A^{\rm ex}$, $A^{\rm twist}$ can shift the wave vector. However, the shift depends on the twist ($\delta N/N$) and the wave vector around the tube axis ($\mu_1 k_1$). Each energy eigenvalue is determined by μ_1 and μ_2, and the conducting electrons with an integer μ_1 ($-N/2 \le \mu_1 \le N/2 - 1$) form an energy band. This means that coupling between $A^{\rm twist}$ and the conducting electron preserves the time-reversal symmetry of the whole system. This is contrasted with the broken time-reversal symmetry by $A^{\rm ex}$.

Persistent currents are defined by differentiating the ground-state energy ($E_0(\Phi)$) with respect to the magnetic flux $\Phi = A^{\rm ex} \cdot T_w$:

$$I_{\rm pc}(\Phi) = -\frac{\partial E_0(\Phi)}{\partial \Phi}. \tag{6}$$

$E_0(\Phi)$ consists of the energy eigenvalue of the valence electrons and has the period of Φ_0. Therefore, $I_{\rm pc}(\Phi)$ also has the period of Φ_0 and is known to be a saw-tooth shape as a function of Φ. However, to calculate the persistent currents in long systems $|T_w| \gg |C_h|$ (or $Q \gg N$), we do not need to sum over the energy eigenvalue of all the valence electrons. We can just calculate the Fermi velocity of each energy band. In terms of Fermi velocity, the amplitude of the persistent current for the μ_1-th energy band is well approximated by [4]

$$I(\mu_1) = \frac{e v_F(\mu_1)}{|T_w|}, \tag{7}$$

where $|T_w|$ is the system length and $v_F(\mu_1)$ denotes the Fermi velocity of the μ_1-th energy band. We have neglected a higher order correction to the amplitude of order of $I(\mu_1)\mathcal{O}(N/Q)$. We fix μ_1 and expand Eq.(5) around the Fermi level (we assume half-filling and set the Fermi level as $E_{\rm F} = 0$) to obtain the energy dispersion relation of the μ_1th energy band as

$$\mathcal{H}_{\mu_1} = v_F(\mu_1) p_2 - \frac{1}{2m(\mu_1)} p_2^2 + \mathcal{O}(p_2^3), \tag{8}$$

where $p_2 (= \mu_2 k_2)$ denotes the momentum along the axis. Notice that $A^{\rm ex} \cdot T_w$ can be included by replacing p_2 with the covariant momentum and Eq.(8) is for $p_2 > 0$. As for $p_2 < 0$, the energy dispersion relation is defined by $-\mathcal{H}_{\mu_1}$ up to $\mathcal{O}(p_2)$. The coefficient of p_2 defines the Fermi velocity and that of p_2^2 the effective-mass if $v_F(\mu_1) = 0$. They are defined respectively as

$$v_F(\mu_1) = 2ta \left| \sin\left(\frac{2\pi\mu_1}{N}\right) \right|, \tag{9}$$

$$\frac{1}{2m(\mu_1)} = ta^2 \cos\left(\frac{2\pi\mu_1}{N}\right). \tag{10}$$

We now consider the persistent currents in an untwisted torus ($\delta N = 0$). In this case, all energy bands yield the same function for the persistent current, that is, a saw-tooth curve as a function of Φ with the same zero points (the flux for which the amplitude of the current vanishes) but with different amplitudes. In fact, the amplitude of the total current is given by a summation of all amplitudes:

$$I_{\rm tot} = \sum_{\mu_1 = -N/2}^{N/2-1} I(\mu_1) = \frac{2eta}{|T_w|} \cot\frac{\pi}{N}. \tag{11}$$

Persistent current $I_{\rm pc}$ in the torus is given by $I_{\rm tot}$ multiplied by ϕ/π ($\phi \equiv 2\pi(\Phi/\Phi_0)$). The linear relation between Φ and the persistent currents has a periodicity of Φ_0, and we then have a saw-tooth current:

$$I_{\rm pc}^{\delta N = 0}(\phi) = I_{\rm tot} \frac{2}{\pi} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{\sin(n\phi)}{n}. \tag{12}$$

The saw-tooth persistent current is a characteristic of the non-interacting theories at zero temperature. Each saw-tooth curve loses its sharpness due to disorder, or at finite temperature [7].

Next, we analyze a twisted torus ($\delta N \neq 0$, see Fig. 1(c)). As we have already shown, the twist behaves as an extra gauge field and shifts the wave vector along the axis direction, so that the zero points of the persistent current also shift [6]. As a result, in order to calculate the total current, we must sum the saw-tooth curves that have different zero points and different amplitudes, both of which depend on μ_1. This can be achieved in a straightforward manner, and the formula for the persistent currents in a twisted torus is given by

$$I_{\rm pc}^{\delta N}(\phi) = \frac{2eta}{|T_w|} I_N^{\delta N}(\phi), \tag{13}$$

where $I_N^{\delta N}(\phi)$ is expressed as

$$
\begin{aligned}
I_N^{\delta N}(\phi) &= \frac{2}{\pi}\sum_{\mu_1=-N/2}^{N/2-1}\sum_{n=1}^{\infty}(-1)^{n+1}\frac{\sin\left(n(\phi-2\pi\mu_1\frac{\delta N}{N})\right)}{n}\left|\sin\left(\frac{2\pi\mu_1}{N}\right)\right| \\
&= \frac{1}{\pi}\sum_{n=1}^{\infty}(-1)^{n+1}\frac{\sin(n\phi)}{n}\left[1+\cos(n\pi\delta N)\right]\left[\frac{2\sin\left(\frac{2\pi}{N}\right)}{\cos\left(\frac{2\pi n\delta N}{N}\right)-\cos\left(\frac{2\pi}{N}\right)}\right]. \qquad (14)
\end{aligned}
$$

Persistent currents exhibit the following characteristics depending on an even or odd number of δN. Term $1+\cos(\pi n\delta N)$ in Eq.(14) vanishes if $n\delta N$ is an odd number or is equal to 2 for other cases. Then Eq.(14) reduces to

$$
I_N^{\delta N}(\phi) = \begin{cases} -\dfrac{1}{\pi}\displaystyle\sum_{n=1}^{\infty}\frac{\sin(2n\phi)}{n}C_n^o & \text{for } \delta N = \text{odd}, \\ \dfrac{2}{\pi}\displaystyle\sum_{n=1}^{\infty}\frac{(-1)^{n+1}\sin(n\phi)}{n}C_n^e & \text{for } \delta N = \text{even}, \end{cases} \qquad (15)
$$

where we have introduced the sequences

$$
C_n^o = \frac{2\sin\left(\frac{2\pi}{N}\right)}{\cos\left(\frac{4\pi n\delta N}{N}\right)-\cos\left(\frac{2\pi}{N}\right)}, \qquad (16)
$$

$$
C_n^e = \frac{2\sin\left(\frac{2\pi}{N}\right)}{\cos\left(\frac{2\pi n\delta N}{N}\right)-\cos\left(\frac{2\pi}{N}\right)}. \qquad (17)
$$

The difference in periodicity for ϕ is clear because $\sin(2n\phi)$ has a period one half the fundamental unit of magnetic flux, or $\Phi_0/2$. This conclusion is valid for any number of N, but a *fundamental* flux periodicity for a current demonstrates more interesting behavior for a specific combination of N and δN.

We consider a particular structure for $\delta N/N = 1/3$ where δN is an even number. In this case, the fundamental period of C_n^e is 3, i.e., $C_{n+3}^e = C_n^e$. When $N \gg 1$, we have $C_1^e = C_2^e \approx -8\pi/3N$ and $C_3^e \approx 2N/\pi$. We then obtain

$$
\begin{aligned}
I_N^{\delta N}(\Phi) &= \frac{2}{\pi}\sum_{n=0}^{\infty}\sum_{m=1}^{3}(-1)^{3n+m+1}\frac{\sin((3n+m)\phi)}{3n+m}C_m^e \\
&= \frac{2}{3\pi}\sum_{n=1}^{\infty}(-1)^{n+1}\frac{\sin(3n\phi)}{n}C_3^e + \mathcal{O}(1/N) \\
&\approx \frac{2N}{\pi^2}\phi \text{ for } N \gg 1, \qquad (18)
\end{aligned}
$$

where in the last line of Eq.(18) we assume $-\pi/3 < \phi < \pi/3$. The fundamental flux period of Eq.(18) becomes $\Phi_0/3$. It should be noted that this argument can be applied to other twisted structures. For instance, when $\delta N/N = 1/Z$ $(1/Z = 1/4, 1/5, \cdots)$, we obtain another fractional period of Φ_0/Z. To our knowledge, this is the first time a fractional periodicity for a twisted boundary condition has been found. We note that the total persistent currents are still saw-tooth curves as expected from the non-interacting theories.

Finally, let us remark that the amplitude of the current exhibits a nontrivial dependence on Φ when $N \to \infty$ with a fixed value of δN. If we divide Eq.(15) by N and take the limit of $N \to \infty$, we then have

$$
\begin{aligned}
&\lim_{N\to\infty}\frac{I_N^{\delta N}(\phi)}{N} \\
&= \begin{cases} -\dfrac{2}{\pi^2}\displaystyle\sum_{n=1}^{\infty}\frac{\sin(2n\phi)}{n}\frac{1}{1-4n^2\delta N^2} & \text{for } \delta N = \text{odd}, \\ \dfrac{4}{\pi^2}\displaystyle\sum_{n=1}^{\infty}(-1)^{n+1}\frac{\sin(n\phi)}{n}\frac{1}{1-n^2\delta N^2} & \text{for } \delta N = \text{even}. \end{cases} \qquad (19)
\end{aligned}
$$

When $\delta N \gg 1$, we sum n in the above equations and obtain

$$
\begin{aligned}
&\lim_{N\to\infty}\frac{I_N^{\delta N}(\phi)}{N} \\
&= \begin{cases} \dfrac{\left(\pi^3-\pi^2 2\phi+(2\phi-\pi)^3\right)}{24\pi^2\delta N^2} & \text{for } \begin{cases} 0 \le \phi \le \pi \\ \delta N = \text{odd}, \end{cases} \\ -\dfrac{\phi\left(\pi^2-\phi^2\right)}{3\pi^2\delta N^2} & \text{for } \begin{cases} -\pi \le \phi \le \pi \\ \delta N = \text{even}, \end{cases} \end{cases} \qquad (20)
\end{aligned}
$$

where we have used the following mathematical formula:

$$
\sum_{n=1}^{\infty}\frac{\sin(n\phi)}{n^3} = \frac{(\pi-\phi)\left[\pi^2-(\phi-\pi)^2\right]}{12} \text{ for } 0 \le \phi \le 2\pi. \qquad (21)
$$

In Fig. 2, we plot the persistent currents of Eq.(20) for an even and odd number of δN as a function of ϕ. It should be noted that the persistent currents are not standard saw-tooth curves even though we are considering non-interacting electrons, and the functional shape of Eq.(20) for δN is an odd number similar to $\sin(2\phi)$ but not identical (see Fig. 2).

Let us summarize our results: (1) A fundamental flux period of the ground state can be generally fractional as Φ_0/Z, depending on the ratio of twist δN to N, although we do not assume any interaction that can form a quasi-particle of charge Ze; (2) As $N \to \infty$ with fixed value $\delta N (\gg 1)$, the currents are not standard saw-tooth as is normally expected from non-interacting theories; and (3) To observe fractional periodicity it is essential that the



FIG. 2: Persistent currents as $N \to \infty$ (Eq.20) for an even (solid line) and an odd (dashed line) number of δN. We multiply 2 (16) for an even (odd) number case. We also plot $0.8\sin(2\phi)$ (dotted line) for comparison.

system contain many electrons: $N \gg 1$ and $Q \gg N$, indicating a kind of "many body" effect.

The primary factor in these results is Fermi surface structure of the square lattice. It appears that many energy bands cross the Fermi level when we roll a planar sheet into a cylinder, and electrons near the Fermi level in each band contribute to the persistent currents, which interfere with one another due to the twist. This point can be made clear by comparing a square lattice with a honeycomb lattice (which possesses only two distinct Fermi points). In this case, we can observe at most a $\Phi_0/2$ periodicity [6].

Here, we discuss some possible extensions of our results. Because we have observed a fractional periodicity in the ground state, one may ask the following question: "Is it possible that the persistent currents show multiple periods such as $2\Phi_0$ or $3\Phi_0$ as a fundamental period?" To answer this question, we consider higher genus materials (g: number of holes), the ground state of which exhibit $g\Phi_0$ periodicity depending on the genus ($g = 2, 3, \cdots$) [8]. A planar system comprising a finite number of square lattices is an example of higher genus material. We note that the conducting electrons in that case are also assumed to be non-interacting and have a single electron charge. Moreover, other examples are known where the charge of a quasiparticle itself becomes fractional. The quasiparticle in the quantum Hall effect has fractional charge $e/3$ [9] and there is a model containing a fractional $e/2$ charged soliton in 1+1 dimensions [10]. Those systems might exhibit $3\Phi_0$ or $2\Phi_0$ periodicity in the ground state respectively. However, fractional periods other than $\Phi_0/2$ (which may correspond to a multiple charge $e \to 3e, 4e, \cdots$) have thus far not been observed as far as we know.

So far we have ignored the dependence of persistent currents on the value of Q and implicitly assumed that Q is a multiple of N. For a general Q, the persistent currents are not so simple [6] as the results obtained in this study. For example, when the reminder of Q/N is an odd number, the one-half periodicity may emerge even in the absence of twist, moreover, another fractional period may appear for a specific value of Q.

In summary, we have examined the flux periodicity of the ground state of the conducting electron on a torus of square lattice. We found that the persistent currents and the ground-state energy of the systems show fractional periods of the fundamental unit of magnetic flux ($\Phi_0/2, \Phi_0/3, \cdots$) depending on twist δN and N. Furthermore, for the case of $N \to \infty$ with a fixed value of $\delta N (\gg 1)$, the persistent currents are not standard saw-tooth curves as expected from the non-interacting theories at zero temperature.

Acknowledgments

K. S. is supported by a fellowship of the 21st Century COE Program of International Center of Research and Education for Materials of Tohoku University. R. S. acknowledges a Grant-in-Aid (No. 13440091) from the Ministry of Education, Japan.

---

[1] Y. Aharonov and D. Bohm, Phys. Rev. **115**, 485 (1959).

[2] M. Büttiker, Y. Imry, and R. Landauer, Phys. Lett. A **96**, 365 (1983); R. Landauer and M. Büttiker, Phys. Rev. Lett. **54**, 2049 (1985).

[3] R.A. Webb, S. Washburn, C.P. Umbach, and R.B. Laibowitz, Phys. Rev. Lett. **54**, 2696 (1985); L.P. Lévy, G. Dolan, J. Dunsmuir, and H. Bouchiat, Phys. Rev. Lett. **64**, 2074 (1990); V. Chandrasekhar, R.A. Webb, M.J. Brady, M.B. Ketchen, W.J. Gallagher, and A. Kleinsasser, Phys. Rev. Lett. **67**, 3578 (1991); D. Mailly, C. Chapelier, and A. Benoit, Phys. Rev. Lett. **70**, 2020 (1993).

[4] Y. Imry, *Introduction to Mesoscopic Physics*, Oxford University Press, New York (1997).

[5] R. Saito, G. Dresselhaus, and M.S. Dresselhaus, *Physical Properties of Carbon Nanotubes*, Imperial College Press, London (1998).

[6] K. Sasaki and Y. Kawazoe, arXiv:cond-mat/0307339.

[7] M. Büttiker, Phys. Rev. B **32**, 1846 (1985); H.F. Cheung, Y. Gefen, E.K. Riedel, and W.H. Shih, Phys. Rev. B **37**, 6050 (1988).

[8] K. Sasaki, Y. Kawazoe, and R. Saito, Phys. Lett. A **321**, 369 (2004).

[9] D.C. Tsui, H.L. Stormer, and A.C. Gossard, Phys. Rev. Lett. **48**, 1559 (1982); R.B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983).

[10] R. Jackiw and C. Rebbi, Phys. Rev. D **13**, 3398 (1976).